A PSEUDO-PARABOLIC VARIATIONAL INEQUALITY AND STEFAN PROBLEM

E. DIBENEDETTO

Mathematics Research Center, University of Wisconsin, Madison, WI 53706, U.S.A.

and

R. E. SHOWALTER

Department of Mathematics, RLM 8.100, The University of Texas, Austin, Texas 78712, U.S.A.

(Received 24 August 1981)

Key Words: Variational inequality, pseudo-parabolic, free-boundary problem,

1. INTRODUCTION

LET A AND B be (possibly multi-valued) maximal monotone operators and let C be a non-empty closed convex set in the real Hilbert space V. We shall give existence and uniqueness results for evolution inequalities (formally) of the form

$$u(t) \in C: \left(\frac{\mathrm{d}}{\mathrm{d}t}(Au(t)) + Bu(t) - f(t), v - u(t)\right)_{V} \ge 0, \qquad v \in C, \quad 0 \le t \le T, \qquad (1.1a)$$

$$(Au(0) - v_0, v - u(0))_v \ge 0, \quad v \in C,$$
 (1.1b)

where $f \in L^2(0, T; V)$ and $v_0 \in A(u_0)$ are given. In section 2 we introduce a new notion of weak solution of (1.1) and verify uniqueness when A is linear self-adjoint and B is strictly monotone. Existence of a weak solution is proved in section 3 when A is a (single-valued) function of the form "identity plus compact operator", B is bounded, and A or B is a subgradient.

Variational inequalities of the form (1.1) are of interest on their own as extensions of corresponding evolution equations of Sobolev type (where C = V). Early work on such inequalities is described in [2]; we mention [6] specifically as a source of examples of initial-boundary-value problems for the pseudo-parabolic partial differential equation

$$\frac{\partial}{\partial t}(u - a\Delta u) = k\Delta u \tag{1.2}$$

with a > 0, k > 0. Such equations arise as models for diffusion, and they provide an interesting alternative to the classical diffusion equation wherein a = 0. In section 4 we give an example of an initial-boundary-value problem consisting of a highly nonlinear partial differential equation of pseudo-parabolic type whose solution is subject to unilateral constraints. Existence and uniqueness results for weak solutions follow from our abstract results on (1.1).

A one-phase free-boundary problem of Stefan type for the equation (1.2) is shown in section 5 to lead to the variational inequality (1.1). This development is parallel to that of the classical case a = 0 which is described, e.g., in [7]. The existence of a classical solution of a Stefan problem for (1.2) in one spatial dimension was given in [9] by entirely different methods.

E. DIBENEDETTO and R. E. SHOWALTER

2. THE VARIATIONAL INEQUALITY

We denote by $L^2(0, T; V)$ the Hilbert space of (Bochner) square-integrable functions on the interval (0, T) with values in the Hilbert space V. Let $H^1(0, T; V)$ denote the absolutely continuous V-valued functions v whose derivatives dv/dt belong to $L^2(0, T; V)$. Denote the dual of V by V^* and recall the natural identification $L^2(0, T; V) = L^2(0, T; V)^*$; thus we obtain the (dual) identification $L^2(0, T; V) \to H^1(0, T; V)^*$ by restriction. The derivative $d/dt : H^1(0, T; V) \hookrightarrow L^2(0, T; V)$ is a bounded linear operator which determines the dual operator $L \equiv -(d/dt)^* : L^2(0, T; V) \to H^1(0, T; V)^*$ by the formula

$$\langle Lf, v \rangle \equiv -\int_0^T (f(t), v'(t))_V dt, \quad f \in L^2(0, T; V), \quad v \in H^1(0, T; V).$$

The restriction of Lf to V-valued test functions is the (distribution) derivative df/dt. Moreover, for $f \in H^1(0, T; V)$ we have

$$\langle Lf, v \rangle = \left(\frac{\mathrm{d}f}{\mathrm{d}t}v\right)_{L^{2}(0, T; V)} + (f(0), v(0))_{V} - (f(T), v(T))_{V}, \quad v \in H^{1}(0, T; V).$$

Thus, we can regard "Lf + f(T)" as formally equivalent to the Cauchy operator "df/dt + f(0)."

We shall use basic material on maximal monotone operators [1]. Specifically, recall $A \subset V \times V$ is monotone if $[x_j, y_j] \in A$ for j = 1 and 2 imply $(x_1 - x_2, y_1 - y_2)_V \ge 0$, and strictly monotone if in addition equality holds only if $x_1 = x_2$. If $\varphi : V \to \mathbf{R} \cup \{+\infty\}$ is proper, convex and lower semicontinuous, its subgradient, defined by

$$\partial \varphi(x) \equiv \{ u \in V : (u, y - x)_V \leq \varphi(y) - \varphi(x) \text{ for all } y \in V \}$$

for $x \in V$, is maximal monotone. More specifically, if C is a non-empty, convex and closed set in V, its indicator function

$$I_{c}(x) \equiv \begin{cases} 0 & , & x \in C \\ +\infty, & x \notin C \end{cases}$$

is proper, convex and lower semicontinuous, and we have $u \in \partial I_{C}(x)$ if and only if

$$x \in C$$
: $(u, y - x) \leq 0$ for all $y \in C$.

Thus, the subgradient of the indicator function provides a convenient method of expressing the variational inequality.

Suppose we are given the pair A, B of maximal monotone operators on the Hilbert space V, a closed convex subset C of V, $f \in L^2(0, T; V)$ and a pair $[u_0, v_0] \in A$. Then a function u is called a *strong solution* of (1.1) if there is a pair of functions v, w such that

$$u, v \in H^{1}(0, T; V); \qquad w \in L^{2}(0, T; V),$$

$$u(t) \in C: \left(\frac{dv(t)}{dt} + w(t) - f(t), x - u(t)\right)_{V} \ge 0, \qquad x \in C,$$

$$v(t) \in A(u(t)) \quad \text{and} \quad w(t) \in B(u(t)) \quad \text{for a.e. } t \in [0, T],$$
(2.1a)

-2

and

$$(v(0) - v_0, x - u(0))_V \ge 0, \quad x \in C.$$
 (2.1b)

Note that since u and v are continuous, C is closed in V and A is closed in $V \times V$, it follows that the inclusions $u(t) \in C$ and $v(t) \in A(u(t))$ hold for all $t \in [0, T]$. Also, (2.1) can be restated as

$$\frac{\mathrm{d}v(t)}{\mathrm{d}t} + w(t) + \partial I_{c}(u(t)) \ni f(t), \qquad (2.2a)$$

 $v(t) \in A(u(t))$ and $w(t) \in B(u(t))$ for a.e. $t \in [0, T]$, and

$$v(0) + \partial I_c(u(0)) \ni v_0 \tag{2.2b}$$

in terms of the indicator function.

We shall use a weak notion of solution in which it is not required that $v \in H^1(0, T; V)$. Set $K \equiv \{u \in H^1(0, T; V) : u(t) \in C, 0 \le t \le T\}$. Define a weak solution of (1.1) to be a function u for which there is a pair of functions v, w satisfying

$$u \in K;$$
 $v, w \in L^{2}(0, T; V),$
 $v(t) \in A(u(t)), w(t) \in B(u(t)),$ a.e. $t \in [0, T],$

and for some $\xi \in A(u(T))$ we have

$$\langle Lv + w - f, \eta - u \rangle + (\xi, \eta(T) - u(T))_V \ge (v_0, \eta(0) - u(0))_V, \quad \eta \in K.$$
 (2.3)

Note that if u is a strong solution then it is a weak solution with $\xi = v(T)$. Moreover we have the following elementary result.

THEOREM 1. Let A be continuous, linear, self-adjoint and monotone; let B be strictly monotone. Then there is at most one weak solution.

Proof. Let u_1 and u_2 be weak solutions and let v_1 , w_1 and v_2 , w_2 be the corresponding selections from $A(u_1)$, $B(u_1)$, etc. By our assumptions on A we have (after modification on a null set) $v_j = A(u_j) \in H^1(0, T; V)$ and $\xi_j = A(u_j(T))$ for j = 1, 2. Thus we have

$$\langle LAu_1 + w_1 - f, u_2 - u_1 \rangle + \langle Au_1(T), u_2(T) - u_1(T) \rangle_V \ge \langle v_0, u_2(0) - u_1(0) \rangle_V$$

 $\langle LAu_2 + w_2 - f, u_1 - u_2 \rangle + (Au_2(T), u_1(T) - u_2(T))_V \ge (v_0, u_1(0) - u_2(0))_V.$

For any $u \in H^1(0, T; V)$ we have

$$\langle LAu, u \rangle = \frac{1}{2} ((Au(0), u(0))_V - (Au(T), u(T))_V)$$

so adding the two inequalities and applying this identity with $u = u_1 - u_2$ gives

$$(w_1 - w_2, u_1 - u_2)_{L^2(0, T; V)} + \frac{1}{2}(Au(T), u(T))_V + \frac{1}{2}(Au(0), u(0))_V \le 0.$$

Strict monotonicity of B shows $u_1 = u_2$.

Remarks. Without additional assumptions we should not expect uniqueness of the selections v, w. For example, in the extreme case $C = \{0\}$, (2.3) is vacuous and we need only choose $v, w \in L^2(0, T; V)$ with $v(t) \in A(0)$ and $w(t) \in B(0)$ to obtain a weak solution. At the other extreme, C = V, any weak solution gives a strong solution of the equation dv/dt + w = f in $L^2(0, T; V)$ with $v(0) = v_0$. Even for equations, the current uniqueness proofs require, e.g., A or B to be linear self-adjoint. See [5].

E. DIBENEDETTO and R. E. SHOWALTER

3. EXISTENCE OF A WEAK SOLUTION

Our objective is to prove the following result on the existence of weak solutions of (1.1). Note that each of our hypotheses concerns only one of the three sources of nonlinearity in the problem; we have not placed any "compatibility" conditions on the operators A, B or the set C.

THEOREM 2. Let C be a non-empty, closed and convex subset of the Hilbert space V. Let A and B be maximal monotone operators on V and assume the following:

- (i) A is a (single-valued) function which maps bounded sets in V into bounded sets in U, where U is a Banach space compactly imbedded in V.
- (ii) B maps bounded sets in V into bounded sets in V.
- (iii) Either $A = \partial \varphi$ or $B = \partial \varphi$, where $\varphi : V \to \mathbf{R}$ is a convex and lower-semicontinuous function.

Then for each $u_0 \in C$ and $f \in L^2(0, T; V)$ there is at least one pair u, w such that

$$u \in K$$
, $w \in L^{2}(0, T; V)$, $w(t) \in B(u(t))$, $0 \le t \le T$, (3.1a)

(3.1b)

$$\langle L(u + A(u)) + w - f, \eta - u \rangle + (u(T) + A(u(T)), \eta(T) - u(T))_V \ge (u_0 + A(u_0), \eta(0) - u(0))_V,$$

for
$$\eta \in K$$
,

and $u(0) = u_0$.

Remarks. If in addition we had $A(u) \in H^1(0, T; V)$, then u would be a strong solution of

$$\frac{d}{dt}(u(t) + A(u(t))) + w(t) + \partial I_c(u(t)) \ni f(t), \quad \text{a.e. } t \in [0, T]$$
(3.2)

This is (2.2a) with A replaced by A + I.

Since we do have $u \in H^1(0, T; V)$ and $u(0) = u_0$, it follows that (3.1b) is equivalent to

$$\left\langle \frac{\mathrm{d}u}{\mathrm{d}t} + LA(u) + w - f, \eta - u \right\rangle + \left(A(u(T)), \eta(T) - u(T) \right)_{V} \ge \left(A(u_{0}), \eta(0) - u(0) \right)_{V}, \qquad \eta \in K.$$
(3.3)

Proof. We shall prove theorem 2 in the following steps. First we approximate (3.2) (and, hence, (3.1)) by replacing ∂I_c by its Lipschitz-continuous Yoshida approximation ∂I_c^{ϵ} , $\epsilon > 0$; the resulting equation has a solution u_{ϵ} by [5]. Then we establish estimates on $\{u_{\epsilon}\}$, deduce the existence of a weak limit $u \equiv \lim(u_{\epsilon})$, and finally show u is a weak solution of (3.1).

The approximation. As an approximation of the indicator function I_c we take

$$I_C^{\varepsilon}(x) \equiv (2\varepsilon)^{-1} \|x - P_C(x)\|_V^2, \qquad \varepsilon > 0, \quad x \in V.$$

Its Fréchet derivative is $\partial I_C^{\epsilon}(x) = \varepsilon^{-1}(x - P_C(x))$, where P_C is the orthogonal projection onto C, and it is monotone and Lipschitz continuous. Thus $B + \partial I_C^{\epsilon}$ is maximal monotone and we obtain from [5] the existence of a pair $u_{\epsilon} \in H^1(0, T; V)$, $w_{\epsilon} \in L^2(0, T; V)$ for each $\varepsilon > 0$ satisfying

$$d/dt \left(u_{\varepsilon}(t) + A(u_{\varepsilon}(t))\right) + w_{\varepsilon}(t) + \partial I_{C}^{\varepsilon}(u_{\varepsilon}(t)) = f(t),$$

$$w_{\varepsilon}(t) \in B(u_{\varepsilon}(t)), \quad \text{a.e. } t \in [0, T],$$

$$(3.4)$$

and $u_{s}(0) = u_{0}$. This approximation (3.4) is strongly suggested by (3.2).

The estimates. Consider the two cases in (iii).

Case $A = \partial \varphi$: Take the scalar product of (3.4) with u_r and integrate; this gives

$$\begin{split} \frac{1}{2} \|u_{\varepsilon}(t)\|_{V}^{2} + \varphi^{*}(A(u_{\varepsilon}(t))) + \int_{0}^{t} (w_{\varepsilon}, u_{\varepsilon})_{V} + \int_{0}^{t} (\partial I_{C}^{\varepsilon}(u_{\varepsilon}), u_{\varepsilon})_{V} \\ &= \frac{1}{2} \|u_{0}\|_{V}^{2} + \varphi^{*}(A(u_{0})) + \int_{0}^{t} (f, u_{\varepsilon}), \qquad 0 \leq t \leq T, \end{split}$$

where $\varphi^*(x) = \sup\{(x, y)_V - \varphi(y) : y \in V\}$ is the convex conjugate of φ [1, p. 41]. Since A is bounded, its domain is all of V so $\varphi(0) < \infty$. Thus, we may take $\varphi(0) = 0$ and $\varphi^*(x) \ge 0$, $x \in V$, with no loss of generality. Since B is monotone $(w_{\epsilon}(t), u_{\epsilon}(t))_V \ge (B^0(0), u_{\epsilon}(t))$ for $t \in [0, T]$, where, e.g., $B^0(0) \in B(0)$ is the minimal section of B at 0. Finally we may assume $0 \in C$ and thus

$$\left(\partial I^{\varepsilon}_{C}(u_{\varepsilon}(t)), u_{\varepsilon}(t)\right)_{V} \geq I^{\varepsilon}_{C}(u_{\varepsilon}(t)) \geq 0, \qquad 0 \leq t \leq T,$$

from the definition of the subgradient. These observations and the preceding estimate give

$$\|u_{\varepsilon}(t)\|_{V}^{2} \leq \frac{1}{2} \|u_{0}\|_{V}^{2} + \varphi^{*}(Au_{0}) + (\|f\|_{L^{1}(0,T;V)} + T\|B^{0}(0)\|_{V}) \|u_{\varepsilon}\|_{L^{\infty}(0,T;V)}, \qquad 0 \leq t \leq T.$$

This implies that $\|u_{\varepsilon}\|_{L^{\infty}(0, T; V)}$ is bounded, and from (i) and (ii) it follows $\|w_{\varepsilon}\|_{L^{\infty}(0, T; V)}$ and $\|A(u_{\varepsilon})\|_{L^{\infty}(0, T; U)}$ are bounded uniformly in $\varepsilon > 0$. Next, we take the scalar product of (3.4) with $u'_{\varepsilon}(t)$ and integrate. This gives

$$\begin{split} \int_{0}^{t} \|u_{\varepsilon}'\|_{V}^{2} + \int_{0}^{t} \left(\frac{\mathrm{d}}{\mathrm{d}t}A(u_{\varepsilon}), \frac{\mathrm{d}u_{\varepsilon}}{\mathrm{d}t}\right)_{V} + I_{C}^{\varepsilon}(u_{\varepsilon}(t)) \\ & \leq I_{C}^{\varepsilon}(u_{0}) + (\|w_{\varepsilon}\|_{L^{2}(0, T; V)} + \|f\|_{L^{2}(0, T; V)}) \|u_{\varepsilon}'\|_{L^{2}(0, T; V)}. \end{split}$$

The monotonicity of A implies the second term above is non-negative so we deduce that $\|u_{\epsilon}'\|_{L^{2}(0,T;V)}$ and $\|I_{C}^{\epsilon}(u_{\epsilon})\|_{L^{\infty}}$ are bounded uniformly in $\epsilon > 0$.

Case $B = \hat{c}\varphi$: Take the scalar product of (3.4) with $u'_{s}(t)$; this gives

$$\|u_{\varepsilon}'(t)\|_{V}^{2} + \left(\frac{\mathrm{d}}{\mathrm{d}t}(Au_{\varepsilon}(t)), \frac{\mathrm{d}}{\mathrm{d}t}u_{\varepsilon}(t)\right)_{V} + \left(w_{\varepsilon}(t) + \partial I_{C}^{\varepsilon}(u_{\varepsilon}(t)), u_{\varepsilon}'(t)\right)_{V} = (f(t), u_{\varepsilon}'(t))_{V}.$$

The second term is non-negative because A is monotone. The third term is the derivative of $\varphi(u_{\varepsilon}(t)) + I_{C}^{\varepsilon}(u_{\varepsilon}(t))$ by the chain rule [1, p. 73]. Thus we integrate this identity and obtain

$$\int_{0}^{\cdot} \|u_{\varepsilon}'\|_{V}^{2} + \varphi(u_{\varepsilon}(t)) + I_{C}^{\varepsilon}(u_{\varepsilon}(t)) \leq \varphi(u_{0}) + \|f\|_{L^{2}(0, T; V)} \|u_{\varepsilon}'\|_{L^{2}(0, T; V)}$$

We can add to $B = \partial \varphi$ a constant, by adding the same to f(t), so we may add an affine function to φ with no loss of generality and thereby obtain $\varphi(x) \ge 0$ for all $x \in V$. The preceding estimate gives uniform bounds on $\|u'_{\varepsilon}\|_{L^{2}(0, T; V)}$ and $\|I^{\varepsilon}_{C}(u_{\varepsilon})\|_{L^{\infty}}$. Similar bounds follow immediately on $\|u_{\varepsilon}\|_{L^{\infty}(0, T; V)}$ and by (i) and (ii) on $\|A(u_{\varepsilon})\|_{L^{\infty}(0, T; V)}$ and $\|w_{\varepsilon}\|_{L^{\infty}(0, T; V)}$, respectively.

The limit. From the estimates obtained above it follows there is a subnet of $\{u_{i}\}$ (which we denote again by $\{u_{i}\}$) for which

$$u - \lim(u_{\varepsilon}) = u \text{ in } H^{1}(0, T; V), \text{ and } w - \lim(w_{\varepsilon}) = w \text{ in } L^{2}(0, T; V),$$

where "w-lim" denotes the weak limit.

w

LEMMA 1. $u \in K \equiv \{v \in H^1(0, T; V) : v(t) \in C, \text{ all } t \in [0, T]\}, u(0) = u_0 \text{ and } w\text{-lim}(u_{\varepsilon}(t)) = u(t) \text{ in } V$ for every $t \in [0, T]$.

Proof. Let t > 0. For each $x \in V$ we have

$$(u_{\varepsilon}(t) - u(t), x)_{V} = \int_{0}^{t} ((u_{\varepsilon} - u)', x)_{V} + (u_{0} - u(0), x)_{V}$$

convergent to $(u_0 - u(0), x)_{\nu}$. By bounded convergence

$$\lim \int_0^T (u_{\varepsilon} - u, x)_V = \int_0^T (u_0 - u(0), x)_V = 0,$$

so $u(0) = u_0$ and w-lim $(u_{\varepsilon}(t)) = u(t)$. Next define $z_{\varepsilon}(t) = P_C(u_{\varepsilon}(t))$, the orthogonal projection onto C. Then $\{z_{\varepsilon}\}$ is bounded in $L^2(0, T; V)$ so there is a subnet $\{z_{\varepsilon}\}$ which converges weakly to z in $L^2(0, T; V)$. Note that

$$\|u_{\varepsilon}(t) - z_{\varepsilon}(t)\|_{V}^{2} = 2\varepsilon I_{C}^{\varepsilon}(u_{\varepsilon}(t)) \leq (\text{const.})\varepsilon,$$

so w-lim $(u_{\xi}) = u = z$. Since the set $\{v \in L^2(0, T; V) : v(t) \in C, \text{ a.e. } t \in C\}$ contains each z_{ξ} and is weakly closed, it also contains u = z. Finally $u \in K$ follows since C is closed and u is continuous.

LEMMA 2. We have the (strong) limits $\lim A(u_{\varepsilon}) = A(u)$ in $L^2(0, T; V)$ and $\lim A(u_{\varepsilon}(t)) = A(u(t))$ in V for every $t \in [0, T]$.

Proof. Let $t \in [0, T]$. Since $\{A(u_{\varepsilon}(t))\}$ is in a compact set in V there is a subnet $\{A(u_{\xi}(t))\}$ which converges (strongly) to v(t) in V. But $w - \lim(u_{\xi}(t)) = u(t)$ in V and A is maximal monotone so v(t) = A(u(t)). The above applies as well to any subnet of $\{A(u_{\varepsilon}(t))\}$, so the entire net converges to A(u(t)). The convergence in $L^2(0, T; V)$ of $\{A(u_{\varepsilon})\}$ to A(u) follows by the bounded convergence theorem.

LEMMA 3. $w \in B(u)$ and $\lim_{\varepsilon} (w_{\varepsilon}, u_{\varepsilon})_{L^{2}(0, T; V)} = (w, u)_{L^{2}(0, T; V)}$.

Proof. It suffices to show that [1, p. 27]

$$\limsup(w_{\varepsilon}, u_{\varepsilon})_{L^{2}(0, T; V)} \leq (w, u)_{L^{2}(0(T; V)}.$$
(3.5)

 (\mathbf{n}, \mathbf{c})

Take the scalar product of (3.4) with $u_{e} - u$ and integrate. From the estimate

$$(\partial I_C^{\varepsilon}(u_{\varepsilon}), u_{\varepsilon} - u)_{L^2(0, T; V)} \ge I_C^{\varepsilon}(u_{\varepsilon}) - I_C(u) = I_C^{\varepsilon}(u_{\varepsilon}) \ge 0$$

we obtain

$$(w_{\varepsilon}, u_{\varepsilon})_{L^2} \leq (w_{\varepsilon}, u)_{L^2} + \left(\frac{\mathrm{d}}{\mathrm{d}t}(u_{\varepsilon} + A(u_{\varepsilon})), u - u_{\varepsilon}\right)_{L^2} + (f, u_{\varepsilon} - u)_{L^2}.$$

By taking the upper limit we have

$$\limsup(w_{\varepsilon}, u_{\varepsilon})_{L^{2}} \leq (w, u)_{L^{2}} + \limsup\left(\frac{\mathrm{d}}{\mathrm{d}t}(u_{\varepsilon} + A(u_{\varepsilon})), u - u_{\varepsilon}\right)_{L^{2}}.$$

Thus it suffices to show the last term is non-positive. From the identity

$$\int_0^T (u_{\varepsilon}', u_{\varepsilon} - u)_V = \frac{1}{2} (\|u_{\varepsilon}(T)\|_V^2 - \|u_0\|_V^2) - \int_0^T (u_{\varepsilon}', u)_V$$

and lemma 1 there follows

$$\liminf \int_0^T (u_{\varepsilon}', u_{\varepsilon} - u)_V \ge \frac{1}{2} (\|u(T)\|_V^2 - \|u_0\|_V^2) - \int_0^T (u', u)_V = 0$$

Similarly from lemma 2

$$\int_0^T (A(u_{\varepsilon})', u_{\varepsilon} - u)_V = (Au_{\varepsilon}(T), u_{\varepsilon}(T) - u(T))_V - \int_0^T (A(u_{\varepsilon}), u_{\varepsilon}' - u')_V$$

it follows that $\lim_{t \to 0} \int_0^T (A(u_{\varepsilon})', u_{\varepsilon} - u)_V = 0.$

The solution. To show that u is a weak solution, it suffices by lemma 1 to verify (3.1b). From any $\eta \in K$ it follows from (3.4) that

$$(u_{\varepsilon}', \eta - u_{\varepsilon})_{L^{2}(0, T; V)} + (A(u_{\varepsilon})', \eta - u_{\varepsilon})_{L^{2}(0, T; V)} + (w_{\varepsilon} - f, \eta - u_{\varepsilon})_{L^{2}(0, T; V)} = (\partial I_{C}^{\varepsilon}(u_{\varepsilon}), u_{\varepsilon} - \eta)_{L^{2}(0, T; V)}.$$
(3.6)

From the definition of subgradient it follows that the right side is greater than $I_C^{\varepsilon}(u_{\varepsilon}) - I_C^{\varepsilon}(\eta) = I_C^{\varepsilon}(u_{\varepsilon})$, hence, non-negative. Consider the first term on the left side. We have from $w - \lim u_{\varepsilon}(T) = u(T)$ that

$$(u', u)_{L^2} = \frac{1}{2} (\|u(T)\|_V^2 - \|u_0\|^2) \le \liminf \frac{1}{2} (\|u_{\varepsilon}(T)\|_V^2 - \|u_0\|_V^2) = \liminf (u'_{\varepsilon}, u_{\varepsilon})_{L^2},$$

so there follows

 $\langle Lu, \eta - u \rangle + (u(T), \eta(T) - u(T))_V - (u_0, \eta(0) - u_0)_V = (u', \eta - u)_{L^2} \ge \limsup(u'_{\epsilon}, \eta - u_{\epsilon})_{L^2}$. Concerning the second term, we obtain from lemma 2

$$\langle L(Au), \eta - u \rangle + (A(u(T)), \eta(T) - u(T))_{V} - (A(u_{0}), \eta(0) - u_{0})_{V}$$

= $\lim \{ \langle L(Au_{\varepsilon}), \eta - u_{\varepsilon} \rangle + (A(u_{\varepsilon}(T)), \eta(T) - u_{\varepsilon}(T))_{V} - (A(u_{0}), \eta(0) - u_{0})_{V} \}$
= $\lim (A(u_{\varepsilon})', \eta - u_{\varepsilon})_{L^{2}(0, T; V)}.$

Finally, lemma 3 identifies the limit of the third term, so by taking the "lim sup" in (3.6) we obtain (3.1.b).

4. A PSEUDO-PARABOLIC INEQUALITY

When our results from above are used to describe initial-boundary-value problems for partial differential equations or inequalities, it is usually more convenient to express them in terms of the equivalent notion of a maximal monotone operator \mathscr{A} from the Hilbert V to its dual V^* . Thus, letting $\mathscr{R}: V \to V^*$ be the Riesz isomorphism given by the scalar product,

$$\mathscr{R}x(y) = (x, y)_V, \qquad x, y \in V,$$

we say $\mathscr{A} \subset V \times V^*$ is monotone if the composite operator $A \equiv \mathscr{R}^{-1} \circ \mathscr{A}$ is monotone in $V \times V$ and maximal monotone if, in addition, $Rg(\mathscr{R} + \mathscr{A}) = V^*$. We can easily state theorem 2, for example, in this context. Thus, we are given a set C closed and convex in $V; \mathscr{A}$ and \mathscr{R} are maximal monotone operators from V to V^* satisfying hypotheses corresponding to (i). (ii) and (iii). Then for each $u_0 \in C$ and $f \in L^2(0, T; V^*)$ there is a pair of functions u, w satisfying

$$u \in K = \{ v \in H^1(0, T; V) : v(t) \in C \text{ for } 0 \le t \le T \},$$
(4.1a)

E. DIBENEDETTO and R. E. SHOWALTER

$$w \in L^2(0, T; V^*), w(t) \in \mathscr{B}(u(t))$$
 for a.e. $t \in [0, T],$ (4.1b)

$$\mathcal{A}(u) \in L^2(0, T; V^*) \text{ and } u(0) = u_0,$$
 (4.1c)

$$\langle L(\mathscr{R} + \mathscr{A})(u) + w - f, \eta - u \rangle + (\mathscr{R} + \mathscr{A})u(T)(\eta(T) - u(T)) \geq (\mathscr{R} + \mathscr{A})u_0(\eta(0) - u(0)), \quad \text{for } \eta \in K.$$

$$(4.1d)$$

In this setting the linear operator $L: L^2(0, T; V^*) \to H^1(0, T; V)^*$ is given by

$$\langle Lg, u \rangle = -\int_0^T g(t)(u'(t)) dt, \quad g \in L^2(0, T; V^*), \quad u \in H^1(0, T; V).$$

Since $u \in H^1(0, T; V)$, the inequality (4.1) is equivalent to

$$\left\langle \frac{\mathrm{d}}{\mathrm{d}t} \,\mathscr{R}u + L\mathscr{A}(u) + w - f, \eta - u \right\rangle + \mathscr{A}(u(T)) \left(\eta(T) - u(T)\right) \ge \mathscr{A}(u_0) \left(\eta(0) - u(0)\right),$$
for $\eta \in K$.

We shall describe an example of a partial differential equation of pseudo-parabolic type which is to be resolved subject to unilateral boundary constraints. (A similar equation with constraint over the entire region will be given in the next section.) Let G be a bounded open set in \mathbb{R}^n which lies on one side of its boundary ∂G ; assume ∂G consists of two disjoint parts Γ_0 and Γ , and let $\overline{n}(s) = (n_1(s), \ldots, n_n(s))$ be the unit outward normal at each point $s \in \partial G$. $H^1(G)$ is the Sobolev space of those $v \in L^2(G)$ for which all derivatives $\partial v/\partial x_j = D_j v$, $1 \le j \le n$, belong to $L^2(G)$; we set $D_0 v = v$. Let $V = \{v \in H^1(G) : v|_{\Gamma_0} = 0\}$; by $v|_{\Gamma_0}$ we mean the trace of v on Γ_0 (see [8, 10]). For $v \in V$ we denote by $\gamma(v) \in L^2(\Gamma)$ the trace of v on Γ . Let $r_0 \in L^{\infty}(G)$ and $r_1 \in L^{\infty}(\Gamma)$ be nonnegative and define

$$\mathscr{R}u(v) = (u, v)_{V} = \int_{G} \left(\sum_{j=1}^{n} D_{j}uD_{j}v + r_{0}uv \right) + \int_{\Gamma} r_{1}\gamma(u)\gamma(v), \qquad u, v \in V.$$

$$(4.2)$$

It follows by a compactness argument that (4.2) is equivalent to the usual $H^1(G)$ scalar-product if any one of Γ_0 or $\{x : r_0(x) > 0\}$ or $\{s : r_1(s) > 0\}$ has strictly positive measure, and we assume this hereafter.

The operator \mathscr{A} is given by a pair of continuous (maximal) monotone functions α_0, α_{-1} : $\mathbf{R} \to \mathbf{R}$ which are linearly bounded:

$$|\alpha_{j}(z)| \leq Q(1+|z|), \quad z \in \mathbf{R}, \quad j = 0, -1$$

for some constant Q > 0. We define

$$\mathscr{A}(u)(v) = \int_{G} \alpha_{0}(u)v + \int_{\Gamma} \alpha_{-1}(\gamma u)\gamma v, \qquad u, v \in V,$$
(4.3)

This operator is a subgradient (in fact, a Gâteaux derivative) and is bounded from V to $U \equiv L^2(G) \times L^2(\Gamma)$. Since the imbeddings $V \hookrightarrow U$ and $U = U^* \hookrightarrow V^*$ are compact, the hypothesis (i) is fulfilled.

The operator \mathscr{B} will be specified by a family of maximal monotone operators $\beta_k : \mathbf{R} \to \mathbf{R}$ which are linearly bounded:

$$|w| \leq Q(1+|z|)$$
 for $w \in \beta_k(z)$, $z \in \mathbf{R}$, $-1 \leq k \leq n$

for some Q > 0. Each $\beta_k = \partial \varphi_k$ for a corresponding convex continuous $\varphi_k : \mathbf{R} \to \mathbf{R}$. We then define $\mathscr{B} = \partial \varphi$ where

$$\varphi(v) \equiv \int_G \sum_{j=0}^n \varphi_j(D_j v) + \int_{\Gamma} \varphi_{-1}(\gamma v), \qquad v \in V.$$

Thus *B* is given (formally) by

$$\mathscr{B}(u) = \sum_{k=0}^{n} D_{k}^{*} \beta_{k}(D_{k}u) + \gamma^{*} \beta_{-1}(\gamma u), \qquad u \in V.$$

$$(4.4)$$

To be precise, we have $F \in \mathscr{B}(u)$ if and only if there exist $f_k \in \beta_k(D_k u)$ in $L^2(G)$, $0 \le k \le n$, and $f_{-1} \in \beta_{-1}(\gamma u)$ in $L^2(\Gamma)$ for which

$$F(v) = \int_G \sum_{k=0}^n f_k D_k v + \int_{\Gamma} f_{-1}(\gamma v), \qquad v \in V.$$

By restricting each of the functionals $\mathscr{R}(u)$, $\mathscr{A}(u)$ and $\mathscr{R}(u)$ to test functions $C_0^{\infty}(G)$ we obtain the corresponding distributions over G

$$\mathscr{R}_0(u) = -\Delta u + r_0 u, \tag{4.5a}$$

$$\mathscr{A}_0(u) = \alpha_0(u), \tag{4.5b}$$

$$\mathscr{B}_{0}(u) = -\sum_{j=1}^{n} D_{j}\beta_{j}(D_{j}u) + \beta_{0}(u), \quad u \in V,$$
 (4.5c)

where the multi-valued \mathscr{B}_0 is interpreted as before. The respective differences are given (by Green's theorem for the first and last cases) for sufficiently regular u by

$$\mathscr{R}u(v) - \int_{\mathcal{G}} \mathscr{R}_{0}(u)v = \int_{\Gamma} \left(\frac{\partial u}{\partial n} + r_{1}\gamma u\right)\gamma v,$$
(4.6a)

$$\mathscr{A}(u)(v) - \int_{G} \mathscr{A}_{0}(u)v = \int_{\Gamma} \alpha_{-1}(\gamma u)\gamma v, \qquad (4.6b)$$

$$\mathscr{B}(u)(v) - \int_{G} \mathscr{B}_{0}(u)v = \int_{\Gamma} \left[\sum_{j=1}^{n} \beta_{j}(D_{j}u)n_{j} + \beta_{-1}(\gamma u) \right] \gamma v, \quad v \in V.$$
(4.6c)

Thus we have realized the operators \mathcal{R} , \mathcal{A} and \mathcal{B} as the sum of a distribution over G (4.5) and a boundary part over Γ (4.6). See [5] for details.

The remaining data is given as follows. Let $C = \{v \in V : \gamma(v) \ge 0 \text{ a.e. on } \Gamma\}$ and let $u_0 \in C$ be specified. Suppose $F_0 \in L^2(G \times [0, T])$ and $g_0 \in L^2(\Gamma \times [0, T])$ are given and define $f \in L^2(0, T; V^*)$ by

$$f(t)(v) = \int_G F_0(.,t)v + \int_{\Gamma} g_0(.,t)\gamma v, \qquad v \in V.$$

With the preceding data as given, the solution u, w of (4.1) is a generalized solution of the pseudo-

parabolic problem

٦

where Λ is the boundary operator obtained from (4.6) as

$$\Lambda(u) = \frac{\partial}{\partial t} \left(\frac{\partial u}{\partial n} + r_1 \gamma(u) + \alpha_{-1}(\gamma u) \right) + \sum_{j=1}^n \beta_j(D_j u) n_j + \beta_{-1}(\gamma u).$$
(4.8)

The other operators in (4.7) are given by (4.5). Note specifically that the multi-valued operators are to be interpreted precisely as was done above following (4.4).

5. A STEFAN PROBLEM

We consider a problem of heat diffusion involving a solid-liquid phase change at a prescribed temperature. One application we have in mind is the melting of ice (initially at temperature zero) suspended in a reservoir or porous medium. The novelty in this treatment is that we assume the heat diffusion is governed by the pair of equations

$$\frac{\partial \theta}{\partial t} = k\Delta \varphi, \qquad \theta = \varphi - a\Delta \varphi.$$

Chen and Gurtin [3] introduced such a model for heat conduction in non-simple materials where the energy, entropy, heat flux and thermodynamic temperature $\theta(x, t)$ depend on the conductive temperature $\varphi(x, t)$ and its first two spatial gradients. Here the heat flux is determined by the conductive temperature and the phase is determined by the thermodynamic temperature. Thus $\theta > 0$ in the region occupied by water and $\theta = 0$ corresponds to the frozen region.

We describe the geometry of the problem. Let the bounded domain G in \mathbb{R}^n be the medium in which the ice/water is suspended and let its boundary ∂G consist of two disjoint pieces, Γ_0 and Γ_1 . Set $\Omega = G \times (0, T)$, where T > 0, and note that its lateral boundary is $B_0 \cup B_1$, where $B_j = \Gamma_j \times (0, T)$ for j = 0, 1. The water-region $\Omega_1 = \{(x, t) \in \Omega : \theta(x, t) > 0\}$ is separated from the ice-region $\Omega_0 = \{(x, t) \in \Omega : \theta(x, t) = 0\}$ by an interface S which is the phase boundary. The unit outward normal on $\partial \Omega_1$ is denoted by $\dot{N} = (\vec{N}_x, N_t), \vec{N}_x \in \mathbb{R}^n$. If V(t) is the velocity in \mathbb{R}^n of the interface at time t, then it follows by the chain rule that $V(t) \cdot \vec{N}_x + N_t = 0$ on S. Set $\vec{n} = \hat{N}_x / \|\vec{N}_x\|$, the unit outward normal in \mathbb{R}^n of the lateral boundary of Ω_1 . Of course $n = \vec{N}_x$ on B_1 , and $\vec{N}_x = 0$ where t = 0 or t = T.

The problem is formulated as follows. We are given the conductivity k > 0, temperature discrepancy a > 0, and latent heat b > 0, of the material and a constant $h \ge 0$ representing conductivity across the lateral boundary B_1 . The initial thermodynamic temperature $\theta_0(x), x \in G$, and applied conductive temperature $g(x, t), (x, t) \in B_1$, are given with $\theta_0 = 0$ on $\Gamma_0, \theta_0 > 0$ on Γ_1 , and $g \ge 0$. The local form of the problem is to find a pair of non-negative functions θ, φ

288

on Ω for which we have

$$\frac{\partial \theta}{\partial t} = k\Delta \varphi, \quad \theta = \varphi - a\Delta \varphi \quad \text{in } \Omega_1$$
 (5.1a)

$$k\frac{\partial\varphi}{\partial n} + bV(t)\cdot\vec{n} = 0$$
 on S (5.1b)

$$k\frac{\partial\varphi}{\partial n} + h(\varphi - g) = 0$$
 on Γ_1 (5.1c)

$$\varphi = 0 \qquad \qquad \text{on } \Gamma_0 \tag{5.1d}$$

$$\theta(.,0) = \theta_0 \qquad \text{on } G. \qquad (5.1.e)$$

Note that if θ , φ is a solution of (5.1) and $\theta_0 \ge 0$, then

$$(a/k)\frac{\partial\theta}{\partial t} + \theta = \varphi \qquad \text{in }\Omega \tag{5.2}$$

so it follows that $\varphi = 0$ on $\Omega_0 \cup S$. Since $g \ge 0$, the maximum principle for the elliptic equation in (5.1a) on the region $G(t) \equiv \{x \in G : (x, t) \in \Omega_1\}$ shows that $\varphi > 0$ in Ω_1 and $\partial \varphi / \partial n < 0$ on S. Thus $N_t < 0$ on S and G(t) is increasing with t.

We shall show that the problem (5.1) leads to a variational inequality of the form (1.1). Define $V = \{v \in H^1(G) : v|_{\Gamma_0} = 0\}$ as before. Regarding regularity of a solution, we assume $\theta_0 \in V$, $\theta : [0, T] \to V$ is absolutely continuous, $\varphi \in L^1(0, T; V)$, and (c.f. (5.2))

$$\frac{a}{k}\frac{\mathrm{d}\theta(t)}{\mathrm{d}t} + \theta(t) = \varphi(t), \qquad \text{a.e. } t \in [0, T].$$
(5.3)

Define the continuous linear $\mathscr{R}: V \to V^*$ by

$$\mathscr{R}u(v) = \int_G k(\nabla u \cdot \nabla v) \,\mathrm{d}x + \int_{\Gamma_1} h(uv) \,\mathrm{d}s, \quad u, v \in V.$$

For a test function $v \in C_0^{\infty}((0, T), V)$ we obtain

$$\int_{0}^{T} \mathscr{R}\varphi(t)(v(t)) dt = \int_{\Omega_{1}} k \nabla \varphi \cdot \nabla v \, dx \, dt + \int_{B_{1}} h \varphi v \, ds \, dt$$
$$= \int_{\Omega_{1}} (-k\Delta\varphi) v \, dx \, dt + \int_{\partial\Omega_{1}} k \nabla \varphi \cdot \overline{N}_{x} v \, ds \, dt + \int_{B_{1}} h \varphi v \, ds \, dt$$
$$= -\int_{\Omega} \frac{\partial \theta}{\partial t} v + \int_{B_{1}} h g v + \int_{S} b N_{t} v$$

from (5.1). Furthermore we have

$$\int_{S} N_{t} v = \int_{\Omega_{1}} \frac{\partial v}{\partial t} = \int_{\Omega} H(\theta) \frac{\partial v}{\partial t} = - \frac{\partial H(\theta)}{\partial t} (v)$$

in the sense of V*-valued distributions, where H(s) = 1 for s > 0 and H(s) = 0 for $s \le 0$ is

the Heaviside function. We can summarize the above calculations as

$$\frac{\mathrm{d}}{\mathrm{d}t}(\theta + bH(\theta)) + \mathscr{R}\varphi = (gh)_{\Gamma_1} \quad \text{in } L^1(0, T; V^*), \tag{5.4}$$

where we define

$$(hg)_{\Gamma_1}(t)(v) = \int_{\Gamma_1} hg(s,t) v(s) \,\mathrm{d}s, \quad v \in V, \quad t \in [0,T].$$

Combining (5.3) and (5.4) we find that the absolutely continuous function $\theta : [0, T] \to V$ satisfies

$$\frac{\mathrm{d}}{\mathrm{d}t}(\theta + (a/k)\mathscr{R}(\theta) + bH(\theta)) + \mathscr{R}(\theta) = (gh)_{\Gamma_1} \quad \text{in } L^1(0, T; V^*), \tag{5.5a}$$

$$\theta(0) = \theta_0, \tag{5.5b}$$

and

$$\theta(x,t) \ge 0, \qquad \text{a.e. } x \in G, \quad t \in [0,T].$$

$$(5.5c)$$

If we integrate (5.5a) and follow the suggestion in [7] to set

$$u(t) \equiv \int_0^t \theta(s) \, \mathrm{d}s$$
$$f(t) \equiv (I + (a/k)\mathcal{R} + bH)\theta_0 - b + \int_0^t (hg)_{\Gamma_1}(s) \, \mathrm{d}s$$

there follows

$$\frac{\mathrm{d}}{\mathrm{d}t}(I+(a/k)\mathscr{R})u+\mathscr{R}u-f(t)=b(1-H(\theta)).$$

Finally we note that $H(u) = H(\theta)$ since G(t) is increasing in t, hence, $u(1 - H(\theta)) = 0$ in Ω . The preceding computations show that $u \in H^1(0, T; V)$ and it satisfies u(0) = 0,

$$u(t) \ge 0 \qquad \text{in } V \tag{5.6a}$$

$$\frac{\mathrm{d}}{\mathrm{d}t}(\mathscr{R} + (k/a)I)u(t) + (k/a)\mathscr{R}u(t) \ge f(t) \quad \text{in } V^*, \tag{5.6b}$$

and

$$\left(\frac{\mathrm{d}}{\mathrm{d}t}(\mathscr{R} + (k/a)I)u(t) + (k/a)\mathscr{R}u(t) - f(t)\right)(u(t)) = 0, \qquad 0 \leq t \leq T.$$
(5.6c)

Setting $C = \{v \in V : v \ge 0 \text{ a.e. in } G\}$ we see that u is a strong solution of (4.1) with $\mathscr{A} = (k/a)I$, $\mathscr{B} = (k/a)\mathscr{R}$, and $u_0 = 0$. Theorem 1 asserts the uniqueness of a solution of (5.1) under conditions considerably weaker than those leading to (5.6). Theorem 2 establishes the existence of a weak solution with certain additional regularity properties. In particular $\mathscr{A}u \in H^1(0, T; V^*)$ since \mathscr{A} is continuous and linear, so (4.1d) is equivalent to

$$\left\langle \frac{\mathrm{d}}{\mathrm{d}t}\left(\mathscr{R}u+\mathscr{A}u\right)+w-f,\eta-u\right\rangle \geq 0, \quad \text{for } \eta\in K.$$

REFERENCES

- 1. BREZIS H., Operateurs Maximaux Monotones et Semigroupes de Contraction dans les Espaces d'Hilbert, Math. Studies 5, North Holland/American Elsevier, (1973).
- 2. CARROLL R. W. & SHOWALTER R. E., Singular and Degenerate Cauchy Problems, vol. 27, Mathematics in Science and Engineering, Academic Press, New York, (1976).
- 3. CHEN P. J. & GURTIN M. E., On a theory of heat conduction involving two temperatures, Z. angew. Math. Phys. 19, 614-627 (1968).
- 4. DIBENEDETTO E. & SHOWALTER R., Stefan problem for pseudo-parabolic equations, in preparation.
- 5. DIBENEDETTO E. & SHOWALTER R. E., Implicit degenerate evolution equations and applications, SIAM J. math. Analysis, 12, 731-751 (1981).
- 6. LAGNESE J., Perturbations in variational inequalities, J. math. Analysis Applic. 55, 302-328 (1976).
- 7. LIONS J. L., Sur Quélques Question d'Analyse de Mechanique et de Contrôl Optimal, Les Presses de l'Universite de Montreal, Montreal (1976).
- 8. LIONS J. L. & MAGENES E., Problemes aux Limites Non Homogenes et Applications, vol. 1, Dunod, Paris, (1967).
- 9. RUNDELL W., The Stefan problem for a pseudo-heat equation, Indiana Univ. Math. J., 27, 739-750 (1976).
- 10. SHOWALTER R. E., Hilbert Space Methods for Partial Differential Equations, Pitman, London (1977).
- 11. TING T. W., A cooling process according to two-temperature theory of heat conduction, J. math. Analysis Applic., 45, 23-31 (1974).